

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/371875902>

Insights on the diet of *Buthus Leach, 1815* (Scorpiones: Buthidae) from Ouarsenis mountains of northwestern Algeria

Article in Revista Ibérica de Aracnología · June 2023

CITATIONS

0

READS

426

5 authors, including:



Chedad Abdelwahab
Université Kasdi Merbah Ouarda

54 PUBLICATIONS 185 CITATIONS

[SEE PROFILE](#)



Mohamed Ait Hammou
Université Ibn Khaldoun Tiaret

86 PUBLICATIONS 607 CITATIONS

[SEE PROFILE](#)



Mayssara El Bouhissi
University of Sidi-Bel-Abbes

26 PUBLICATIONS 39 CITATIONS

[SEE PROFILE](#)



Salah Eddine Sadine
Université de Ghardaïa

88 PUBLICATIONS 349 CITATIONS

[SEE PROFILE](#)

Insights on the diet of *Buthus* Leach, 1815 (Scorpiones: Buthidae) from Ouarsenis mountains of northwestern Algeria

Abdelwahab Chedad^{1,2*}, Mohamed Ait Hammou³, Mayssara El Bouhissi^{4,5},
Abdessalam Chedad⁶, & Salah Eddine Sadine⁷

¹ Department of Biological Sciences, Laboratory of Saharan Bio-Resources: Preservation and Valorization, Ouargla University,.30000 Ouargla, Algeria. Tel./fax.: +213-778-610632, agrochedad@yahoo.fr, ORCID: <https://orcid.org/0000-0001-8098-1803>

² Directorate of Forest Conservation of Ghardaïa. (General Directorate of Forestry), 47000 Ghardaïa, Algeria.

³ Department and Faculty of Nature and Life Sciences, Laboratory of Agro Biotechnology and Nutrition in Semi-Arid Areas, Ibn Khaldoun University of Tiaret. 14000 Tiaret, Algeria. mohamedaithammou@gmail.com

⁴ Laboratory Ecodevelopment of spaces, Djilali Liabes University. Sidi Bel Abbes, Algeria. elbouhissimayssara@yahoo.fr⁵ Directorate of Forest Conservation of Sidi Bel Abbes, (General Directorate of Forestry), 22000 Sidi Bel Abbes, Algeria

⁶ Independent Naturalist, Bordj Bounaama. Tissemsilt, 38001, Algeria, abdelwahabchedad@gmail.com.

⁷ Laboratoire Valorisation et Conservation des Ecosystèmes Arides (LVCEA). Faculté des Sciences de la Nature et de la Vie et Sciences de la terre, Université de Ghardaïa, Ghardaïa, Algeria. sse.scorpion@yahoo.fr

Abstract: This study is an empirical observation of the feeding behavior of the *Buthus* Leach, 1815, in the wild of the Ouarsenis Mountains (Tissemsilt and Tiaret regions, northwestern Algeria) during the study period (2021–2022). We have observed 29 instances of *Buthus* species eating arthropods. The identifications of this prey yielded a list of 20 species belonging to 15 families and 9 orders. We note a dominance of the Orthoptera and Coleoptera, with 3 families and 5 species for each. Gryllidae and Scarabaeidae are the most represented, with 3 and 2 species, respectively. Also, we record some insects larvae eating and many cases of cannibalism and intraguild predation. All of the *Buthus* species' prey are intermediate or smaller arthropods.

Key words: Algeria, *Buthus*, Cannibalism, Diet, intraguild, Ouarsenis.

Nota Perspectivas sobre la dieta de *Buthus* leach, 1815 (Scorpiones: Buthidae) de las montañas Ouarsenis del norte de Argelia

Resumen: Este estudio es una observación empírica del comportamiento alimentario de *Buthus* Leach, 1815, en la naturaleza de las montañas Ouarsenis (regiones de Tissemsilt y Tiaret, norte de Argelia) durante el período de estudio (2021–2022). Hemos observado 29 casos de especies de *Buthus* que comen artrópodos. Las identificaciones de esta presa arrojan una lista de 20 especies pertenecientes a 15 familias y 9 órdenes. Observamos una dominancia de Ortópteros y Coleópteros, con 3 familias y 5 especies para cada uno. Gryllidae y Scarabaeidae son los más representados, con 3 y 2 especies, respectivamente. Además, registramos el consumo de larvas de insectos y muchos casos de canibalismo y depredación intragrupo. Todas las presas de la especie *Buthus* son artrópodos intermedios o más pequeños.

Palabras clave: Argelia, *Buthus*, Canibalismo, Dieta, Ouarsenis.

Introduction

Certainly, habitat (microhabitat) confirms foraging success for animals due to prey availability or foraging efficiency (Griffiths, 1975; Werner *et al.*, 1981). Scorpions are solitary, sedentary arthropods that are ecologically important because they influence the flow of energy in arid and semi-arid ecosystems (Polis and McCormick, 1986; Polis, 2001; Brown and Kotler, 2004; Nime *et al.*, 2016). They are efficient predators and accept a wide variety of prey such as invertebrates, small lizards, snakes, and even small mice (Vachon, 1952; Kock, 1969; Polis and McCormick, 1986; McCormick and Polis, 1990; McCormick and Polis, 1995; Teruel, 2015; Lira *et al.*, 2016; Dupré, 2015a; Sadine, 2018; Sadine and El Bouhissi, 2021; Chedad *et al.*, 2022a).

Moreover, they are stationary in nature; thus, they are often encountered in their shelter, which they only leave to feed and reproduce (Polis, 1990). Generally, during the day, they seek shelter under rocks and logs, in crevices and burrows dug in the substrate, or under the loose outer layers of shrubs and trees of many plants. Just after sunset, individuals emerge from these diurnal retreats to forage or engage in other activities (Polis, 1979; Polis, 1990).

Where scorpions enter the diet of many animals, especially vertebrates, Polis *et al.* (1981) summarized a list of approximately 150 taxa of scorpion predators, and Dupré (2015b) resumed a list of 312 species that can be considered scorpion predators, with more than 80% being vertebrates (birds, mammals, and reptiles). Among the few consumers is the wild boar, *Sus scrofa*, whose consumption of scorpions was recorded in the Ouarsenis region (west Algeria) (Chedad *et al.*, 2021).

The genus *Buthus* Leach, 1815, belongs to the most abundant and widely distributed scorpion family, Buthidae C.L. Koch, 1837, with 78 validly named species (Rein, 2022). The number of confirmed *Buthus* species in Algeria has increased to 11: two species have a wide distribution, i.e., *B. paris* and *B. tunetanus*; there are also eight endemic species for Algeria such as *B. tassili* Lourenço, 2002; *B. pusillus* Lourenço, 2013; *B. saharicus* Sadine, Bissati and Lourenço, 2016; *B. aures* Lourenço and Sadine, 2016; *B. boussaadi* Lourenço, Chichi and Sadine, 2018 and *B. apiatus* Lourenço, El Bouhissi and Sadine, 2020; *B. ahaggar* Ythier, Sadine, Haddadi and Lourenço, 2021; *B. goyffoni* Abidi, Sadine and Lourenço, 2021; and a Moroccan species, *B. oudjanii* Lourenço, 2017.

This study aims to provide a contribution to the diet of the *Buthus* species from the Ouarsenis Mountains (northwestern Algeria), based on an empirical observation in the wild environment that includes various events of cannibalism and intraguild predation.

Material and methods

Study area

Our research area is in the Ouarsenis massif, in the Tissemsilt and Tiaret regions (northwestern Algeria) (Fig. 1). This forest massif is characterized by a semi-arid to humid climate at altitudes between 400 and 1985 m, several biotopes with distinct plant covers, and a slope of up to 50% (BNEDER, 2009; Chedad *et al.*, 2022b).




Fig. 1. Map of Algeria, showing the study area (Tissemsilt and Tiaret regions).

For two years (2021–2022), the onset of high temperatures coincides with the beginning of the exit of scorpions from hibernation (the end of April until the end of October). We searched for scorpions with an average rate of once every 10 days, during predation under stones and straws, in burrows and crevices, and we also searched for food scraps, on which we concentrated our outings, especially in the early morning (two to three hours) and the period before sunset (one hour).

The study area (Ouarsenis massif) exhibits a very high level of scorpion-specific diversity, with more than 62% of the species being *Buthus* (Chedad *et al.*, 2022b). The following table summarizes the *Buthus* species that can be found in this region (Table I).

Results and discussion

During the study period (2021–2022), in the wild of the Ouarsenis region (Tissemsilt and Tiaret), we observed 29 cases of *Buthus* species eating arthropods. Morphological identification of this prey yielded a list of 20 species belonging to 15 families and nine orders (Table II).

Our results show the dominance of the Orthoptera and the Coleoptera, with three families and five species for each. Gryllidae and Scarabaeidae are the most represented, with three and two species, respectively (Fig. 1, Tab. II). *Buthus* sp. 1 has the highest abundance (13.79%), while *Messor* sp. has the second highest (10.34%). We also recorded some insects larvae, which are considered fast to digest.

One Sample T-Test ($df = 19,0; p < 0,001$): The p-value for the test is $< 0,001$, which is less than the alpha = 0,05 significance level. It can be concluded that the consumption by *Buthus* species varies according to the type of prey. (fig. 2)

Scorpions are characterized by an external digestion that remains very slow (Quinlan *et al.*, 1995). Their diet is very diverse and tends to be insectivorous. It contains important arthropods abundant in their biotope, such as insects (small beetles, butterflies, locusts, grasshoppers, and ants), crustaceans (woodlice), and arachnids (scorpions, spiders, opiliones, etc.) (Williams, 1987; Sánchez Piñero *et al.*, 2013; Sadine, 2018; Toprak *et al.*, 2022). In our study, the *Buthus* spp. diet is composed of more than 75% of insects from its biotope. However, the scorpions consumed represent more than 20.5% of *Buthus*' diet (Fig. 3). Nevertheless, *Buthus* species, like other scorpions, can feed on small vertebrates (reptiles and rodents) (McCormick & Polis, 1990; Sadine & El Bouhissi, 2021).

Many authors reported that the foraging success of scorpions may be associated with seasonal changes in prey availability (Polis 1980; Polis & McCormick, 1986). In general, it can be summarized into two types of prey capture strategies. First, most species hunt prey by waiting for passages near or just in front of their burrows/hiding places. The second is to actively hunt away from hiding places. (McCormick & Polis, 1990).

Table I. *Buthus* species than can be found in the Ouarsenis region.

No.	Species	References
1	<i>B. apiatus</i> Lourenço, El Bouhissi and Sadine, 2020	Benali and Feghouli, 2022; Chedad <i>et al.</i> , 2022b
2	<i>B. aures</i> Lourenço and Sadine 2016	Benali and Feghouli, 2022; Chedad <i>et al.</i> , 2022b
3	<i>B. boussaadi</i> Lourenço, Chichi and Sadine, 2018	Benali and Feghouli, 2022
4	<i>B. oudjani</i> Lourenço, 2017	Benali and Feghouli, 2022
5	<i>B. paris</i> (C.L. Koch, 1839)	Benali and Feghouli, 2022; Chedad <i>et al.</i> , 2022b
6	<i>B. tunetanus</i> (Herbst, 1800)	Benali and Feghouli, 2022; Chedad <i>et al.</i> , 2022b
7	<i>Buthus</i> sp.	Chedad <i>et al.</i> , 2022b

Table II. Prey of the *Buthus* species in the Ouarsenis region, Algeria.

No.	Orders	Families	Species	Relative abundance (RA%)
01	Araneae	Lycosidae	<i>Hogna</i> sp.	3.45
		Tenebrionidae	<i>Pimelia</i> sp.	6.90
			<i>Scaurus</i> sp.	3.45
02	Coleoptera	Scarabaeidae	<i>Oxythyrea</i> sp.	3.45
			<i>Amphimallon</i> sp.	3.45
03	Dermaptera	Staphylinidae	<i>Ocypus</i> sp.	3.45
		Forficulidae	<i>Forficula</i> sp.	3.45
04	Hymenoptera	Apidae	<i>Xylocopa</i> sp.	3.45
05	Isopoda	Formicidae	<i>Messor</i> sp.	10.34
06		Porcellionidae	<i>Porcellio</i> sp.	3.45
07	Mantodea	Mantidae	<i>Mantis religiosa</i>	6.90
		Gryllidae	<i>Gryllus</i> sp.	6.90
08	Orthoptera	Acrididae	<i>Sciobia</i> sp. 1	6.90
		Tettigoniidae	<i>Sciobia</i> sp. 2	3.45
09	Scolopendromorpha	Scolopendridae	<i>Schistocerca</i> sp.	3.45
			<i>Tettigonia</i> sp.	3.45
			<i>Scolopendra cingulata</i>	3.45
		Buthidae	<i>Buthus</i> sp. 1	13.79
Total	09	Scorpionidae	<i>Buthus</i> sp. 2	3.45
			<i>Scorpio</i> sp.	3.45
			20	100




Fig. 2. Distribution of families of arthropod species identified as prey of *Buthus* spp. in the Ouarsenis region, Algeria.



Fig. 3. Prey of *Buthus* species from Algeria. **A:** *Mantis religiosa*, **B:** *Gryllus* sp., **C:** Larva 1, **D:** Larva 2, **E:** *Scolopendra cingulata*, **F:** *Schistocerca* sp., **G:** *Porcellio* sp., **H:** *Ocypus* sp.



Fig. 4. **A.** Cases of cannibalism *Buthus/Buthus*; **B.** Case of intraguild predation *Buthus/Scorpio*

This study shows cases of cannibalism in *Buthus/Buthus* and intraguild predation in *Buthus/Scorpio* (Fig. 4), where scorpions are considered very voracious predators and cannibals. For some predators, such as scorpions, cannibalism is an important factor in mortality that determines ecology and behavior (Sánchez-Piñero and Urbano-Tenorio, 2016), and it is usually the key interaction that shapes the distribution, behavior, and population structure of these species (Polis, 1981; Leonardsson, 1991; Rudolf, 2006). Predator threat therefore becomes a key factor in determining where animals can safely forage and influencing trophic interactions, population, and community structure (Hawlena and Schmitz, 2010; Cresswell and Quinn, 2013). This phenomenon is an asymmetrical interaction, with larger predators usually feeding on smaller conspecifics (Polis, 1981); this is what we have noted through this study and has also been reported by Chedad et al. (2022a).

Conclusions

This work constitutes the first study that shows an overview of the trophic diet of the *Buthus* species in Northwestern Algeria, where we have observed a remarkable diversity of the trophic menu of these species. Therefore, it becomes necessary to conduct a deep study in this field in order to learn more about the trophic diet and preferences of the scorpions and generalize them to other regions. As scorpions are nocturnal species, it is desirable to monitor their feeding behavior carefully, whether to follow them in captivity or in the wild.

Acknowledgments

The authors thank all accompanying persons especially Ms. Belli M., Boughalem D., Chedad AK., Adli D., Amalou D., Amalou B., Belghazi D., Saad Y. and Waheb R. We also thank the Algerian Ministry of Higher Education, and reviewers of this paper.

References

- ABIDI, H., SADINE, S. E., HOUHAMDI, M., MADOUI, A. & W.R. LOURENÇO 2021. The genus *Buthus* Leach, 1815 in Algeria (Scorpiones: Buthidae) and a possible new case of vicariant species. *Revista Ibérica de Aracnología*, **38**: 81-86.*
- BENALI, N. & M.A., FEGHOUL 2022. *Inventaire, morphologie et distribution des Aracnides de la région de Tiraet*. Thèse de Master en Sciences de la nature et de la vie. Université Ibn Khaldoun-Tiaret. Algérie, 108 pp.
- BNEDER, 2009. Plan national de Développement Forestier (PNDF) (National Forest Development Plan). Rapport de synthèse nationale, Alger, Algérie.
- BROWN, J.S. & B.P. KOTLER 2004. Hazardous duty pay and the foraging cost of predation. *Ecol Lett*, **7**: 999–1014.
- CHEDAD, A., AIT HAMMOU, M. & S.E. SADINE 2021. The wild boar *Sus scrofa* Linnaeus, 1758 as a predator of scorpions (Scorpiones) in north-western Algeria (Mammalia, Suidae). *Revista Ibérica de Aracnología*, **39**: 137–138.*
- CHEDAD, A., AIT HAMMOU, M., CHEDAD, A., LACHICHI, A., BELL M. & S.E. SADINE 2022a. Note on a natural case of *Scorpio maurus* cannibalism in northwestern Algeria (Scorpiones: Scorpionidae). *Revista Ibérica de Aracnología*, **41**: 131-133.*
- CHEDAD, A., AIT HAMMOU, M., CHELGHOU, H., CHEDAD, A., OULD AMARA, O., EL BOUHSSI, M., DAHMANI W. & S.E. SADINE 2022b Diversity and distribution pattern of scorpions from the Ouarsenis massif of Tissemsilt, North-West Algeria. *Biodiversitas*, **23**(5): 2444-2450. <https://doi.org/10.13057/biodiv/d230523>
- CRESSWELL, W. & J.L. QUINN 2013. Contrasting risks from different predators change the overall nonlethal effects of predation risk. *Behav. Ecol.*, **24**: 871–876.
- DUPRÉ, G. 2015a. Les proies des scorpions « Synthèse ». *Arachnidés*, **76**:22-31.
- DUPRÉ, G. 2015b. Les prédateurs des scorpions (Arachnida: Scorpiones). *Arachnidés*, **76** :1-31.
- GRIFFITHS, D. 1975. Prey availability and the food of predators. *Ecology*, **56**:1209–1214.
- HAWLENA, D. & O.J. SCHMITZ 2010. Herbivore physiological response to predation risk and implications for ecosystem nutrient dynamics. *PNAS*, **107**: 15503–15507. doi: 10.1073/pnas.1009300107 PMID:20713698
- HERBST, J.F.W. 1800. Naturgeschichte der Skorpione. *Natursystem der Ungeflügelten Insekten*. Berlin: Bei Gottlieb August Lange, 86 pp.
- KOCH, C.L. 1839. Die Arachniden. C. H. Zeh'sche Buchhandlung. Nürnberg, 6(1-6): 1-156.
- KOCK, D. 1969. Ein Skorpion frist an einer Zwergraus. *Natur Museum Frankfurt*, **99**(3): 107-112.
- LEONARDSSON, K. 1991. Effects of cannibalism and alternative prey on population dynamics of *Saduria entomon* (Isopoda). *Ecology*, **72**: 1273–1285.
- LIRA, A.F.A.; FOERSTER S.I.A. & A.A.C. SILVA-FILHO 2016. Reports of scorpion predation by spiders in the Brazilian Atlantic forest and Caatinga (Arachnida: Scorpiones, Araneae). *Revista Ibérica de Aracnología*, **28**: 87-89.*
- LOURENÇO, W.R. 2002. Considérations sur les modèles de distribution et différentiation du genre *Buthus* Leach, 1815, avec la description d'une nouvelle espèce des montagnes du Tassili des Ajjer, Algérie (Scorpiones, Buthidae). *Biogeographica*, **78**(3): 109-127.
- LOURENÇO, W.R. 2013. A new species of *Buthus* Leach, 1815 from Algeria (Scorpiones, Buthidae). *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, **16**: 63-68.
- LOURENÇO, W.R. AND SADINE, S.E. 2016. One more new species of *Buthus* Leach, 1815 from Algeria (Scorpiones: Buthidae). *Revista Ibérica de Aracnología*, **28**: 13-17.*
- LOURENÇO, 2017 : « Encore une nouvelle espèce de *Buthus* Leach, 1815 (Scorpiones: Buthidae) pour le nord-est du Maroc. » *Revista Ibérica de Aracnología*, **31** : 59-63.*
- LOURENÇO, W.R., CHICHI, S. & S.E. SADINE 2018. A new species of *Buthus* Leach, 1815 from the region of Bou Sâada-M'Sila, Algeria; a possible case of vicariance for the genus (Scorpiones: Buthidae). *Revista Ibérica de Aracnología*, **32**: 15-20.*
- LOURENÇO, W.R., EL BOUHSSI, M. & S.E. SADINE 2020. Further considerations on the *Buthus* Leach, 1815 species present in Algeria with description of a new species (Scorpiones: Buthidae). *Revista Ibérica de Aracnología*, **36**: 103-108.*
- MCCORMICK, S.J. & G.A. POLIS 1990. Prey, predators, and parasites. pp. 294-320 in G.A. Polis (Ed.). *The biology of scorpions*. Stanford, Stanford University Press, 587 pp.
- MCCORMICK, S.G. & G.A. POLIS 1995. Prey, Predators, Parasites. In «The Biology of Scorpions», Polis G. A., Eds., Stanford University Press, Stanford, CA, 294-320.
- NIME, M.F., CASANOVES, F. & C.I. MATTONI 2016. Microhabitat use and behavior differ across sex-age classes in the scorpion *Brachistosternus ferrugineus* (Scorpiones: Bothriuridae). *Journal of Arachnology*, **44**:235–244.
- POLIS GA. 1979. Prey and feeding phenology of the desert sand scorpion *Paruroctonus mesaensis* (Scorpionidae: Vaejovidae). *Journal of Zoology (London)*, **188**:333–346.
- POLIS, G.A. 1981. Evolution and dynamics of intraspecific predation. *Annu. Rev. Ecol. Syst.* **12**: 225–251.
- POLIS, G.A., SISSOM, W.D. & S.J. MCCORMICK 1981. Predators of scorpions: field data and a review. *Journal of Arid Environments*, **4**: 309-326. [https://doi.org/10.1016/s0140-1963\(18\)31477-0](https://doi.org/10.1016/s0140-1963(18)31477-0)
- POLIS, G.A. & S.J. MCCORMICK 1986. Patterns of resource use and age structure among species of desert scorpion. *Journal of Animal Ecology*, **55**:59–74. <https://doi.org/10.2307/4692>
- POLIS, G.A. 1990. In *The biology of scorpions*. (G.A. Polis, ed.). Stanford University Press, Stanford, California. *Ecology*, 247–293.
- POLIS, G.A. 2001. Population and community ecology of desert

- scorpions, pp. 302-316. In: P.H. Brownell & G.A. Polis (Eds.). *Scorpion biology and research*. Oxford, Oxford University Press, 595 pp.
- QUINLAN, T.G., SMITH, G.T. & M.C. CALVER 1995. Relationships between morphology and feeding behaviour in the syntopic scorpion *Urodacus armatus* Pocock and *Urodacus novae-hollandiae* Peters (Scorpions: Scorpionidae). *J. Austr. Ent. Soc.*, **34**: 277-279.
- REIN, J.O. 2020. The Scorpion Files. <https://www.ntnu.no/ub/scorpion-files/> (accessed on 26.12. 2022).
- RUDOLF, V.H.W. 2006. The influence of size-specific indirect interactions in predator-prey systems. *Ecology*, **87**: 362-371. PMID: 16637362
- SADINE, S.E., BISSATI, S. & W.R. LOURENÇO 2016. The first true deserticolous species of *Buthus* Leach, 1815 from Algeria (Scorpiones, Buthidae); ecological and biogeographic considerations. *Comptes Rendus Biologies*, **339**: 44-49. <http://dx.doi.org/10.1016/j.crvi.2015.10.002>
- SADINE, S.E. 2018. La faune scorpionique du Sahara septentrional algérien: Diversité et Ecologie. Thèse de Doctorat ès Sciences. Université Kasdi Merbah-Ouargla. Algérie, 112 pp.
- SADINE, S.E. & M. EL BOUHSSI 2021. A case of predation on *Acanthodactylus* Wiegmann, 1834 (Squamata, Lacertidae) by *Androctonus amoreuxi* (Scorpiones, Buthidae) in western Algeria. *Revista Ibérica de Aracnología*, **39**: 130.*
- SÁNCHEZ PIÑERO, F., URBANO TENORIO, F. & F. JESÚS MARTÍN GARCÍA 2013. Foraging of *Buthus occitanus* (Scorpiones: Buthidae) on shrub branches in an arid area of southeastern Spain," *The Journal of Arachnology*, **41**(1): 88-90. <https://doi.org/10.1636/H11-90.1>
- SÁNCHEZ-PIÑERO, F. & F. URBANO-TENORIO 2016. Watch Out for Your Neighbor: Climbing onto Shrubs Is Related to Risk of Cannibalism in the Scorpion *Buthus cf. occitanus*. *PLoS ONE*, **11**(9):e0161747. <https://doi.org/10.1371/journal.pone.0161747>
- TERUEL, R. 2015. A scorpion predating upon a solifuge in Mexico (Arachnida: Scorpiones, Solifugae). *Revista Ibérica de Aracnología*, **26**: 93-94.*
- TOPRAK, S., KURT, R. & E.A. YAĞMUR 2022. First report of intra-guild predation in scorpions (Scorpiones: Buthidae) from Turkey. *Euscorpius*, **363**: 1-4.
- VACHON, M. 1952. Etudes sur les scorpions. Publications de l'Institut Pasteur d'Algérie, Alger. 482 pp.
- WERNER, E.E., MITTELBACH, G.G. & D.J. HALL 1981. The role of foraging profitability and experience in habitat use by the bluegill sunfish. *Ecology*, **62**:116-125.
- WILLIAMS, S.C. 1987. Scorpion bionomics. *Ann. Rev. Ent.*, **32**: 275-295.
- YTHIER, E., SADINE, S.E., HADDADI, M.L. & W.R. LOURENÇO 2021. « A new species of *Buthus* Leach, 1815 from Algeria (Scorpiones: Buthidae) and an interesting new case of vicariance. » *Faunitaxys*, **9**(21):1-9.

* Reference available at www.sea-enomologi.org